Claims

5

10

15

25

30

- 1. Method of producing a product protein, wherein the protein is expressed from a mammalian cell, preferably from a lymphoid cell, in cell culture at least during a certain span of time during cell culture, comprising the steps
 - e) preparing a cell culture medium for culturing mammalian cells, preferably preparing a cell culture medium that is devoid of butyrate, and further preferred preparing a cell culture medium allowing for growth of the mammalian cells, more preferably a protein-free cell culture growth medium,
 - f) and further adding acetic acid or an acetate salt or an acetyl ester to a final concentration of from 1 to 20 mM, preferably of from 3 to 15 mM, more preferably of from 5 to 12 mM, most preferably of from 6 to 9.5 mM, said addition being carried out either directly to the medium prior to starting cell culture or feeding it to the medium during cell culture,
 - g) further culturing, preferably growing, said cell in said medium with concomittant expression of product protein,
 - h) and finally harvesting said protein from the cell culture.
- 20 2. Method according to claim 1, characterised in the the addition of acetic acid or a salt thereof is carried out directly to the medium prior to or at starting the cell culture.
 - 3. Method according to one of the preceding claims, characterised in that an acetate alkali metal or alkaline earth metal salt is added to the medium.
 - 4. Method according to one of the preceding claims, characterised in that the cells are lymphoid cells, preferably NS0 cells.
 - 5. Method according to claim 4, characterised in the cells are NS0 cells that are recombinant for and can express Glutamine synthetase.
 - 6. Cell culture medium for animal cell culture, characterised in that the medium is suited for culturing mammalian cells and comprises acetic acid or an acetate salt or a

WO 03/064630 PCT/EP03/03631

biologically activated acetyl ester at a concentration of from 1 to 20 mM, preferably of from 3 to 15 mM, more preferably of from 5 to 12 mM, most preferably at about 6 to 9.5 mM, and preferably is devoid of butyric acid or any of its salts.

- 7. Cell culture medium according to claim 3, characterised in that the medium is a high density cell culture medium.
 - 8. Cell culture medium according to claim 6 or 7, characterised in that the medium is a cell culture medium allowing of growth of animal cells, preferably lymphoid cells.
 - 9. Cell culture medium according to claims 6-8, characterised in that the medium is a serum-free and protein-free cell culture medium, preferably a protein-free medium suitable for NS0 cell culture.

10

- 10. A medium concentrate for preparation of a culture medium as defined in claim 6 which is either a solid or a liquid.
 - 11. Use of a cell culture medium according to claim 6 for lymphoid cell culture.
- 12. Cell culture, comprising mammalian cells, preferably lymphoid cells, most preferably NS0 cells, in a medium according to claim 6.